

MDS

Supports Methylation Reactions in the Liver



PRODUCT BENEFITS*

- Supports methylation reactions associated with liver detoxification
- Promotes ideal homocysteine metabolism
- Supports the production and metabolism of bile
- Supports a healthy liver

MDS is a broad-spectrum nutritional formulation that supports methylation reactions and, in turn, supports proper homocysteine metabolism. MDS also supports the production and metabolism of bile as part of healthy liver function.

DIRECTIONS FOR USE

Take 2 capsules once per day, or as directed by your health care professional.

STORAGE

Store in a cool, dry place away from direct sunlight. Product from unopened containers that have been stored under recommended conditions (55-85° F / 13-29° C) may be used for up to 36 months.

WARNING

Consult your health care professional before use if pregnant, nursing, taking medications, or for any use by minors. Do not use if safety seal is broken or missing. Keep out of reach of children.



SUPPLEMENT FACTS

	Amount Per Serving	% Daily Value
Serving size: 2 Capsules		
Servings per container: 45		
Vitamin C (as Ascorbic Acid)	37 mg	41%
Vitamin E (as D-Alpha Tocopheryl Succinate)	8.7 mg	58%
Riboflavin (as Riboflavin and Riboflavin-5-Phosphate Sodium)	7 mg	538%
Niacin (from Niacinamide)	4 mg NE	25%
Vitamin B6 (as Pyridoxal-5-Phosphate)	6 mg	353%
Folate (as L-5-Methyltetrahydrofolate Calcium)	570 mcg DFE	144%
Vitamin B12 (as Methylcobalamin)	505 mcg	21,042%
Vitamin B7 (Biotin)	800 mcg	2,553%
Choline (as Choline Bitartrate)	16 mg	3%
Magnesium (as Magnesium Citrate)	88 mg	21%
Betaine Anhydrous	387 mg	**
Methylsulfonylmethane (MSM)	36 mg	**
Betaine HCl	36 mg	**
SAME (as S-Adenosyl-L-Methionine Disulfate P-Tosylate)	31 mg	**
N-Acetyl-L-Cysteine	15 mg	**

* Percent Daily Values are based on a 2000 calorie diet.

** Daily Value Not Established

Other Ingredients: Gelatin (Capsule), Dicalcium Phosphate, Magnesium Stearate (Vegetable), Silica Hydrophilic Fumed Powder, Pruv® (Sodium Stearyl Fumarate).

*To be provided by health care professionals only. The dosage recommendations are only for your health care provider's consideration. Please consult your health care provider for your individual dosing instructions. This product is for nutritional purposes only. It is not designed to diagnose, treat, reverse, cure, or prevent any disease. This product is not intended to replace or delay the use of prescription medication. These statements have not been evaluated by the FDA. All rights reserved © Biogenetix, LLC

REFERENCE LIST

1. Allen RH, et al. Serum betaine, N,N-dimethylglycine and N-methylglycine levels in patients with cobalamin deficiency and related inborn errors of metabolism. *Metab.* 1993;42(11):1448-1460.
2. Brattstoerm L, et al. Plasma homocysteine in women on oral oestrogencontaining contraceptives and in men with oestrogen-treated prostatic carcinoma. *Scand J Clin Lab Invest.* 52, 1982:283-287.
3. Butterworth CE, et al. Folate deficiency and cervical dysplasia. *JAMA.* 1992; 267:528-533.
4. Finkelstein JD. Homocysteine: a history in progress. *Nutr Rev.* 2000;58(7): 193-204.
5. Krumdieck CL, Prince CW. Mechanisms of homocysteine toxicity on tissues: implications of morbidity of aging. *J Nutr.* 2000;130:365S-368S.
6. Lindenbaum J, et al. Neuropsychiatric disorders caused by cobalamin deficiency in the absence of anemia or macrocytosis. *N Eng J Med.* 1988;318:1720-1728.
7. Smithells RW, et al. Possible prevention of neural-tube defects by periconceptional vitamin supplementation. *Lancet*, Feb 16, 1980:339-342.
8. Smythies JR. The role of the one-carbon cycle in neuropsychiatric disease. *Biol Psychiatry.* 1984;19(5):755-758.
9. Zingh JM, Jones PA. Genetic and epigenetic aspects of DNA methylation on genome expression, evolution, mutation and carcinogenesis. *Carcinogenesis.* 1997; 18(5):869-882.