

BioG-Max Multi+

Ultimate Multi-Vitamin Blend



PRODUCT BENEFITS

- **Protects Against Oxidative Damage**
- **Stimulates Metabolism**
- **Supports Healthy Physiology**

BioG-Max Multi+ is the state-of-the-art multivitamin blend that offers the ultimate in absorption and efficacy. Multi+ packs active forms of B vitamins along with a comprehensive blend of stable and metabolically active vitamins and antioxidants.

Unlike standard blends in the marketplace, Biogenetix utilizes the most absorbable, stable and metabolically active forms of all of our basic vitamins. The Multi+ formula includes highly bioactive B vitamins, Vitamin D3 for maximum effect, both forms of Vitamin K, and trimethylglycine to support the liver and methylation. Tocotrienols, part of the vitamin E family, offer unusually potent antioxidant activity, and three specialized carotenoids can broaden that antioxidant protection, especially safeguarding the eye and vision.



SUPPLEMENT FACTS

	Amount Per Serving	% Daily Value
Serving size: 1 tsp (5 mL)		
Servings per container: 20		
Vitamin A (as Retinol, Beta Carotene)	2100 mcg RAE	233%
Vitamin C (as Sodium Ascorbate)	100 mg	111%
Vitamin D (as Cholecalciferol) (D3)	62.5 mcg (2500 IU)	313%
Vitamin E (as Natural Mixed Tocopherols, Tocofersolan)	23 mg	153%
Vitamin K (as K2 (MK7), K1 (Phytonadione))	90 mcg	75%
Thiamin (Vitamin B1) (Thiamine HCl)	12 mg	1000%
Riboflavin (as Riboflavin-5-Phosphate)	3.6 mg	277%
Niacin (as Niacin, Niacinamide)	10 mg	63%
Vitamin B6 (as Pyridoxine HCl)	6.7 mg	394%
Folate (as Calcium Folate)	850 mcg DFE	213%
Vitamin B12 (as Methylcobalamin)	500 mcg	20833%
Biotin	500 mcg	1667%
Pantothenic Acid (as Calcium d-Pantothenate)	22.5 mg	450%
Sodium (as Sodium Ascorbate)	10mg	<1%
Trimethylglycine (as betaine)	20 mg	**
Proprietary Blend	363 mg	**
Phospholipids (from purified sunflower seed lecithin), liquid Milk Thistle seed extract, Tocotrienols (from annatto), Lycopene, Zeaxanthin, Lutein purified sunflower seed lecithin)		

* Daily Value not established

Other Ingredients: Water, glycerin, ethanol, EDTA (as preservative), natural citrus oils and natural flavoring.

Does Not Contain: Sugar, wheat, gluten, milk, soy, egg, artificial flavorings, colorings, peanuts, tree nuts or fish.

KEY INGREDIENTS

Antioxidants. Daily antioxidant intake has never been more important with the current standard American diet. By providing exquisite support to the body as it fights against oxidative stress, tissue breakdown, and metabolic disease, Multi+ plus promotes optimal levels of inflammation and healthy aging.

Water-Soluble Vitamins. Required in frequent, small doses, these vitamins support the body as it metabolizes macronutrients in energy metabolism, creates and repairs DNA, and develops and maintains new cells and tissue. The water soluble profile in Multi+ consists of the most bioavailable form of each vitamin.

Fat-Soluble Vitamins. To provide phenomenal support to the body as it promotes optimal immune function, healthy membrane and tissue integrity, and ideal hormone balance, Multi+ houses potent bioactive forms of each fat soluble vitamin.

To be provided by health care professionals only. The dosage recommendations are only for your health care provider's consideration. Please consult your health care provider for your individual dosing instructions. This product is for nutritional purposes only. It is not designed to diagnose, treat, reverse, cure, or prevent any disease. This product is not intended to replace or delay the use of prescription medication. These statements have not been evaluated by the FDA. All rights reserved © Biogenetix, LLC

www.biogenetix.com
orders@biogenetix.com
(833) 525-0001

DIRECTIONS FOR USE

Take 1 tsp once daily directly by mouth, hold for 30-60 seconds and swallow. Repeat to desired dosage or as directed by a health care professional. Best taken on an empty stomach at least 10 minutes before meals. If pregnant, breastfeeding, or planning to become pregnant, consult your physician before use.

STORAGE

Store at room temperature.

REFERENCE LIST

1. Schurgers LJ, et al. Blood. 2007;109:3279-83.
2. Peh HY, et al. Pharmacol Ther. 2016;162:152-69.
3. Sen CK, et al. Life Sci. 2006;78:2088-98.
4. Wu W, et al. Mol Nutr Food Res. 2015;59:1663-73.
5. Zou X, et al. Cell Death Dis. 2014;5:e1218.
6. Yang PM, et al. Life Sci. 2016;155:94-101.
7. Alyautdin R, et al. Int J Nanomedicine. 2014;9:795-811.
8. Ahn H, Park JH. Biomater Res. 2016;20:36.
9. Spector AA, Yorek MA. J Lipid Res. 1985;26:1015-35.