

Glucostatic Balance

Supports Healthy Insulin and Blood Sugar Levels



PRODUCT BENEFITS*

- Supports healthy metabolism
- Supports healthy glucose levels
- Promotes healthy insulin levels

Glucostatic Balance is a vitamin, mineral, and herbal based formulation designed to support healthy blood sugar metabolism, normal insulin receptor site sensitivity, and healthy insulin responses. This product also helps to buffer the bodies response to the glycemic load.



DIRECTIONS FOR USE

Take 2 capsules once per day, or as directed by your health care professional.

STORAGE

Store in a cool, dry place away from direct sunlight. Product from unopened containers that have been stored under recommended conditions (55-85° F / 13-29° C) may be used for up to 36 months.

WARNING

Consult your health care professional before use if pregnant, nursing, taking medications, or for any use by minors. Do not use if safety seal is broken or missing. Keep out of reach of children.

SUPPLEMENT FACTS

	Amount Per Serving	% Daily Value
Serving size: 2 Capsules Servings per container: 90		
Vitamin E (as D-Alpha Tocopheryl Succinate)	42.88 mg	286%
Thiamin (as Thiamin HCl)	7mg	583%
Riboflavin	7mg	538%
Niacin (as Niacinamide and Nicotinic Acid)	32 mg	200%
Vitamin B6 (as Pyridoxine HCl)	4 mg	235%
Vitamin B12 (as Cyanocobalamin)	7 mcg	292%
Biotin	428 mcg	1,427%
Pantothenic Acid (as Calcium D-Pantothenate)	43 mg	860%
Choline (as Choline Bitartrate)	57 mg	10%
Magnesium (as Magnesium Citrate)	36 mg	9%
Zinc (as Zinc Glycinate)	7 mg	64%
Selenium (as Selenium Glycinate)	29 mcg	53%
Manganese Bisglycinate Chelate	4 mg	174%
Chromium (as Chromium Polynicotinate)	200 mcg	571%
Proprietary Blend	487 mg	**
N-Acetyl L-Cysteine, Gymnema Leaf Extract, Berberine HCl, R-Alpha Lipoic Acid, L-Carnitine (as L-Carnitine L-Tartrate), Para-Aminobenzoic Acid (PABA), Inositol, Garcinia Fruit Powder, Lagerstroemia speciosa (Banaba) Leaf Powder, Bitter Melon Fruit Powder, Organic Cassia Bark Powder, Fenugreek Seed Powder, Guggul Gum Resin Powder		
* Percent Daily Values are based on a 2000 calorie diet.		
** Daily Value Not Established		

Other Ingredients: Gelatin (Capsule), Cellulose, Silica, Magnesium Stearate.

*To be provided by health care professionals only. The dosage recommendations are only for your health care provider's consideration. Please consult your health care provider for your individual dosing instructions. This product is for nutritional purposes only. It is not designed to diagnose, treat, reverse, cure, or prevent any disease. This product is not intended to replace or delay the use of prescription medication. These statements have not been evaluated by the FDA. All rights reserved © Biogenetix, LLC

REFERENCE LIST

- Abdel-Aziz MT, Abdou MS, Soliman K, et al. Effects of L-carnitine on blood lipid patterns in diabetic patients. *Nutr Rep Int* 1984;29:1071-1079.
- Anderson HU, Jorgensen KH, Egeberg J. Nicotinamide prevents interleukin-1 effects on accumulated insulin release and nitric oxide production in rat islets of langerhans. *Diabetes* 1994;43:770-777.
- Anderson R, Cheng N, Bryden, et al. Beneficial effects of chromium for people with type II diabetes. *Diabetes*. 1996;45:124A.
- Anderson RA, Cheng N, Bryden NA, et al. Elevated intakes of supplemental chromium improve glucose and insulin variables in individuals with type 2 diabetes. *Diabetes* 1997;46:1786-1791.
- Anderson RA, Polansky MM, Bryden NA, Canary JJ. Supplemental-chromium effects on glucose, insulin, glucagon, and urinary chromium losses in subjects consuming controlled low-chromium diets. *Am J Clin Nutr* 1991 ;54:909-916.
- Barbagallo M, Dominquez LJ, Tagliamonte MR, et al. Effects of vitamin E and glutathione on glucose metabolism: role of magnesium. *Hypertension* 1999; 34:1002-1006.
- Boden G, Chen X, Ruiz J, et al. Effects of vanadyl sulfate on carbohydrate and lipid metabolism in patients with non-insulin dependent diabetes mellitus. *Metabolism*. 1996;45:1130-1135.
- Chen MD, Lin PY, Lin Wh. Investigation of relationship between zinc and obesity. *Kao Hsiung I Hsueh Tsa Chih* 1991 ;7:628-634.
- Clearly JP. Vitamin B3 in the treatment of diabetes mellitus: case reports and review of the literature. *J Nutr Med* 1990; 1 :217-225.
- Cohen N, Halberstam M, Shlimovich P, et al. Oral vanadyl sulfate improves hepatic and peripheral insulin insensitivity in patients with non-insulin dependent diabetes mellitus. *J Clin Invest*. 1995;95:2501-2509.
- Dominguez LJ, Barbagallo M, Sowers JR, Resnick LM. Magnesium responsiveness to insulin and insulin-like growth factor I in erythrocytes from normotensive and hypertensive subjects. *J Clin Endocrinol Metab* 1998;83:4402-4407.
- Evans GW, Browman TD. Chromium picolinate increases membrane and rate of insulin internalization. *J Inorg Bio* 1992;46:243-250.
- French RJ, Jones PJ. Role of vanadium in nutrition: metabolism, essential and dietary considerations. *Life Sciences*. 1992;52:339-346.
- Goldfine AB, Simonson DC, Folli F, et al. Metabolic effects of sodium metavanadate in humans with insulin-dependent and noninsulin-dependent diabetes mellitus in vivo and vitro studies. *J Clin Enco and Meta*. 1995;80(11);3311-3319.
- Greco AV, et al. Effect of propionyl-L-carnitine in the treatment of angiopathy: Controlled double blind trial versus placebo. *Drugs Exp Clin Res* 1992;18:69-80.
- Gunal AI, Celiker H, Donder E, Gunal SY. The effect of L-carnitine on insulin resistance in hemodialysed patients with chronic renal failure. *J Nephrol* 1999; 12:38-40.
- Halberstam M, Cohen N, Shlimovich P, et al. Oral vanadyl sulfate improves insulin sensitivity NIDDM but not in obese nondiabetic subjects. *Diabetes*. 1996; 45:659-665.
- Hegazi SM. Effects of zinc supplementation on serum glucose, insulin, glucagons, glucose-6-phosphate, and mineral levels in diabetics. *J Clin Biochem Nutr* 1992;12:209-215.
- Heller W, Mushi HE, Gaebel G, et al. Effect of L-carnitine on post-stress metabolism in surgical patients. *Infusioether Klin Ernahr* 1986;13:268-276.
- Humphries S, Kushner H, Falkner B. Low dietary magnesium is associated with insulin resistance in a sampling of young, nondiabetic Black Americans. *Am J Hypertens* 1999;12:747-756.
- Jacob S, Henriksen EJ, Schieman AL, et al. Enhancement of glucose disposal in patients with Type 2 diabetes by alpha-lipoic acid. *Arzneim-Rosch Drug Res* 1995;45(2):872-874.
- Jacob S, Ruus P, Hermann R, et al. Improvement of insulin-stimulated glucose disposal in type 2 diabetes after repeated patenteral administration of thioctic acid. *Exp Clin Endocrinol Diabetes* 1996;104:284-288.
- Jacob S, Ruus, Herman R, et al. Oral administration of RAC-alpha-lipoic acid modulates insulin sensitivity in patients with type-2 diabetes: a placebocontrolled pilot trial. *Free Radie Biol Med* 1999;27:309-314.
- Jacob S, Streeper RS, Fogt DL, et al. The antioxidant alpha-lipoic acid enhances insulin-stimulated glucose metabolism in insulin resistant rate skeletal muscle. *Diabetes* 1996;45:1024-1029.
- Jain SK, McVie R, Jaramillo JJ, et al. Effect of modest vitamin E supplementation on blood glycated hemoglobin and triglyceride levels and red cell indices in type I diabetic patients. *J Amer Col Nutr* 1996;15(5):458-461.
- Kaskaran K, et al. Antidiabetic effect of a leaf extract from *Gynema sylvestre* in non-insulin dependent diabetes mellitus patients. *J Ethnopharmacol* 1990; 30: 295-305.
- Kawabata T and Packer L. Alpha-lipoate can protect against glycation of serum albumin, bu nto low density lipoprotein. *Biochem Biophys Res Comm* 1994; 203:99-104.
- Khare AK, Tondon RN, Tewari JP. Hypoglycemic activity of an indigenous drug (*Gynema sylvestre*, "Gurmar") in normal and diabetic persons. *Indian J Physiol Pharmacol* 1983;27(3):257-8.
- Konard T, Vicini P, Kuster K, et al. Alpha-lipoic acid treatment decreases serum lactate and pyruvate concentrations and improves glucose effectiveness in lean and obese patients with type 2 diabetes. *Diabetes Care* 1999;22(2):280-287.
- Kutoskos D, Fournounas C, Kepetanaki A, et al. Oral glucose tolerance test after high-dose i.v. biotin administration in normoglycemic hemodialysis patients. *Ren Fail* 1996;18:131-137.
- Leslie ROG, Elliot RB. Early environmental events as a cause of IDDM. *Diabetes* 1994;43:843-850.
- Levine R, Streeten D, Doisy R. Effect of oral chromium supplementation on the glucose tolerance of elderly human subjects. *Metabolism* 1968;17:114-125.
- Maebashi M, Makino Y, Furukawa Y, et al. Therapeutic evaluation of the effect of biotin on hyperglycemia in patients with non-insulin dependent diabetes mellitus. *J Clin Biochem Nutr* 1993;14:211-218.
- Mingrone G, Greco AV, Capristo E, et al. L-carnitine improves glucose disposal in type 2 diabetic patients. *J Am Coll Nutr* 1999;18:77-82.
- Mooradian AD, Morley JE. Micronutrient status in diabetes mellitus. *Am J Clin Nutr* 1987;45:877-895.
- Nadler JL, Buchanan T, Natarajan R, et al. Magnesium deficiency produces insulin resistance and increased thromboxane synthesis. *Hypertension* 1993;21:1024-1029.
- Nagamtsu M, et al. Lipoic acid improves nerve blood flow, reduces oxidative stress, and improves distal nerve conduction in experimental diabetic neuropathy. *Diabetes Care* 1995;18:1160-1167.
- Offenbacher EG, Pi-Sunyer FX. Beneficial effect of chromium-rich yeast on glucose tolerance and blood lipids in elderly subjects. *Diabetes* 1980;29:919-925.
- Paolisso G, D'Amore A, Galzenano D. Daily vitamin E supplements improve metabolic control but not insulin secretion in elderly type II diabetic patients. *Diabetes Care* 1993;16:1433-1437.
- Paolisso G, D'Amore A, Guigliano D, et al. Pharmacologic doses of vitamin E improves insulin action in healthy subjects and in non-insulin-dependent diabetic patients. *Am J Clin Nutr* 1993;57:650-656.
- Paolisso G, Gambardella A, Giugliano D, et al. Chronic intake of pharmacological doses of vitamin E might be useful in the therapy of elderly patients with coronary heart disease. *Am J Clin Nutr* 1995;61 :848-852.
- Paolisso G, Ravussin E. Intercellular magnesium and insulin resistance; results in Pima Indians and Caucasians. *J Endocrinol Metab* 1995;80:1382-1385.
- Paolisso G, Sgambato S, Gambardella A, et al. Daily magnesium supplements improve glucose handling in elderly subjects. *Am J Clin Nutr* 1992;55:1161-1167.
- Paolisso G, Sgambato S, Piza G, et al. Improved insulin response and action by chronic manganese administration in aged NIDDM subjects. *Diabetes Care* 1989;12:265-269.
- Persaud SJ, Al-Majed H, Raman A, Jones PM. *Gynema sylvestre* stimulates insulin release in vitro by increased membrane permeability. *J Endocrinol* 1999; 163(2):207-12.
- Pocoit F, Reimers JI, Anderson HU. Nicotinamide - biological action and therapeutic potential in diabetes prevention. *Diabetologia* 1993;36:574-576.
- Pozzilli P, Andreani D. The potential role of nicotinamide in the secondary prevention of IDDM. *Diabetes Metabol Rev* 1993;9:219-230.
- Reddi A, DeAngelis B, Frank O, et al. Biotin Supplementation improves glucose and insulin tolerances in genetically diabetic KK mice. *Life Science* 1998;42:1323-1330.
- Riales R, Albrink MJ. Effect of chromium chloride supplementation on glucose tolerance and serum lipids including high-density lipoprotein of adult men. *Am J Clin Nutr* 1981 ;34:2670-2678.
- Roebach, Jr. JR, Hla KM, Chambless, et al. Effects of chromium supplementation on serum high density lipoprotein cholesterol levels in men taking beta-blockers. *Ann Intern Med*. 1991 ;115:917-924.
- Rudich A, Tirosh A, Potashnik R, Khamaisi M, Bashan N. Lipoic acid protects against oxidative stress induced impairment in insulin stimulation of protein kinase B and glucose transport in 3T3-L 1 adipocytes. *Diabetologia* 1999;42:949-957.
- Salonen JT, Jyssonen K, Tuominen TP. Increased risk of non-insulin diabetes mellitus at low plasma vitamin E concentrations. A four year followup study in men. *Br Med J* 1995;311 :1124-1127.
- Shang H, Osada K, Maebashi M, et al. A high biotin diet improves the impaired glucose tolerance of long-term spontaneously hyperglycemic rats with non-insulin dependent diabetes mellitus. *J Nutr Sci Vitamin* 1996;42:517-526.
- Shang H, Osada K, Maebashi M, et al. A high biotin diet improves the impaired glucose tolerance of long-term spontaneously hyperglycemic rats with non-insulin-dependent diabetes mellitus. *J Nutr Sci Vitamin* 1996;42:517-526.
- Shanmugasundaram ERB, et al. Possible regeneration of the islet of Langerhans in streptozotocin-diabetic rats given *Gynema sylvestre* leaf extracts. *J Ethnopharmacol* 1990 30; 265-279.
- Shanmugasundaram ERB, et al. Use of *Gynema sylvestre* leaf extract in the control of blood glucose in insulin-dependent diabetes mellitus. *J Ethnopharmacol* 1990; 30: 281-294.
- Shanmugasundaram KR, et al. Enzyme changes and glucose utilization in diabetic rabbits: the effect of *Gynema sylvestre*, R.Br. *J Ethnopharmacol* 1983; 7(2):205-34.
- Shechter, Li Meyerovitch, et al. Insulin-like actions of vanadate are mediated in an insulin receptor independent manner via non-receptor protein tyrosine kinases and protein phosphotyrosine phosphates. *Mol Cell Biochem*. 1995; 153(1-2):39-47.
- Singh RB, Niaz MA, Rastogi SS, et al. Current zinc intake and risk of diabetes and coronary artery disease and factors associated with insulin resistance in rural and urban populations of North India. *J Am Col Nutr* 1998;17:564-570.
- Streeper RS, Henriksen EJ, Jacob S, et al. Differential effects of lipoic acid stereoisomers on glucose metabolism in insulin resistant skeletal muscle. *Am J Physiol* 1997;273:E185-E191.
- Striffler JS, Polansky MM, Anderson RA. Overproduction of insulin in the chromium-deficient rat. *Metabolism* 1999;48:1063-1068.
- Striffler JS, Polansky MM, et al. Chromium improves insulin response to glucose in rats. *Metabolism* 1995;44:1314-1320.
- Urberg M, Zemel MB. Evidence of synergism between chromium and nicotinic acid in the control of glucose tolerance in elderly humans. *Metabolism* 1987; 36:574-576.

*To be provided by health care professionals only. The dosage recommendations are only for your health care provider's consideration. Please consult your health care provider for your individual dosing instructions. This product is for nutritional purposes only. It is not designed to diagnose, treat, reverse, cure, or prevent any disease. This product is not intended to replace or delay the use of prescription medication. These statements have not been evaluated by the FDA. All rights reserved © Biogenetix, LLC

www.biogenetix.com
orders@biogenetix.com
 (833) 525-0001