

# Hepato-ST

Supports Healthy Liver Function



## PRODUCT BENEFITS\*

- Supports healthy liver function and detoxification
- Support the synthesis of phospholipids and lipid metabolism

The purpose of Hepato-ST is four-fold: 1) to support healthy liver function by helping normal distribution of fat in the liver, 2) to support the synthesis of phospholipids, which can help facilitate the elimination of lipids from the liver, and 3) to support liver function.

Hepato-ST contains nutritional compounds that have lipotropic properties and can be useful in the support of a healthy liver and, in particular, support the elimination of lipids from the liver.

Non-alcoholic fatty liver disease (NAFLD) is the most common chronic liver disease in the world. A large contributor to the pathogenesis of NAFLD is chronic underconsumption of choline based on both functional and genetic factors. Choline has shown promise in its ability to support the healthy maintenance of liver tissue and function in those affected by NAFLD.

In addition to the powerful support offered by Choline, Hepato-ST also supplies progressive amounts of broccoli powder, shown to support healthy function of and protect against fatty liver changes.

Classic use of Milk Thistle Extract and herbal promotion of drainage combine with the powerful effects offered by the anti-NAFLD components blend together herbal tradition and cutting-edge application.



## SUPPLEMENT FACTS

Serving size: 1 Capsule Servings per container: 120	Amount Per Serving	% Daily Value
Vitamin B6 (as pyridoxal 5'-phosphate)	2.5 mg	147%
Folate (As L-Methylfolate Calcium) (70 mcg L-Methylfolate)	119 DFE	30%
Choline (as Choline Citrate)	100 mg	18%
Magnesium (as Magnesium Citrate)	5 mg	1%
L-Carnitine (as L-Carnitine L-Tartrate)	125 mg	**
Taurine	30 mg	**
Milk Thistle Seed Extract (Silybum marianum)	30 mg	**
N-Acetyl-L-Cysteine	25 mg	**
Broccoli (Brassica oleracea)(flore)†	25 mg	**
Inositol	25 mg	**
L-Methionine	25 mg	**

\* Percent Daily Values are based on a 2000 calorie diet.

\*\* Daily Value Not Established

**Other Ingredients:** Gelatin (Capsule), Microcrystalline Cellulose, Magnesium Stearate.

\*To be provided by health care professionals only. The dosage recommendations are only for your health care provider's consideration. Please consult your health care provider for your individual dosing instructions. This product is for nutritional purposes only. It is not designed to diagnose, treat, reverse, cure, or prevent any disease. This product is not intended to replace or delay the use of prescription medication. These statements have not been evaluated by the FDA. All rights reserved © Biogenetix, LLC

## DIRECTIONS FOR USE

Take 1 capsule 3 times per day, or as directed by your health care professional.

## STORAGE

Store in a cool, dry place away from direct sunlight. Product from unopened containers that have been stored under recommended conditions (55-85° F / 13-29° C) may be used for up to 36 months.

## WARNING

Consult your health care professional before use if pregnant, nursing, taking medications, or for any use by minors. Do not use if safety seal is broken or missing. Keep out of reach of children.

## REFERENCE LIST

- Adzet T. Polyphenolic compounds with biological activity and pharmacological activity. *Herbs Spices Medicinal Plants* 1986;1:167-184
- Belcaro GV, Grimaldi R, Guidi G. Improvement of capillary permeability in patients with venous hypertension after treatment with TTFCA. *Angiology* 1990; 41(7): 533-540.
- Bohm K. Choleric action of some medicinal plants. *Arzneimittel Forsch* 1959;9: 376-378.
- Bombardelli E, Cirstoni A, Lietta A. The effect of acute and chronic (Panax) ginseng saponins treatment on adrenal function; biochemical and pharmacological. *Proceedings 3rd International Ginseng Symposium*. Seoul: Korean Research Institute. 1980: p 9-16.
- Damis F, Orcel L, de Saint-Maur PP, Mamaou P. Use of a titrated extract of *Centella asiatica* in chronic hepatic disorders. *Sem Hosp Paris* 1979;55:1749-1750.
- El Zawahry MD, Kahil AM, El Banna MH. Madecassol. A new therapy for hepatic fibrosis. *Bull Soc Int Chir (Belgium)* 1975;34:296-297.
- Faber K. The dandelion *Taraxacum officinale*. *Pharmaze* 1958;13:423-436.
- Feibrich F, Koch H. Silymarin an inhibitor of lipoxygenase. *Experientia* 1979;35: 148-150.
- Feibrich F, Koch H. Silymarin, an inhibitor of prostaglandin synthesis. *Experientia* 1979;35:150-152.
- Heafield MT, et al. Xenobiotic metabolism in Alzheimer's disease. *Neurology*. 1990;40:1095-1098.
- Hikino H, Kiso Y, Sandah S, Shoji J. Antihepatotoxic actions of ginsenosides from Panax ginseng roots. *Planta Medica* 1985;52: 62-64.
- Hikino H, Kiso Y, Wagner H. Antihepatotoxic actions of flavanone glycosides from Silybum marianum fruits. *Planta Medica* 1984;50:248-250.
- Jain A, Buist NR, Keenaaway NG, et al. Effect of ascorbate or N-acetylcysteine treatments in patient with hereditary glutathione synthetase deficiency. *J Pediatr* 1994; 124: 229-233.
- Johnston CJ, Meyer CG, Srilakshmi JC. Vitamin C elevates red blood cell glutathione levels in healthy adults. *Am J Clin Nutr* 1993; 58: 103-105.
- Larbovic D. Single blood test might predict drugs' effects on patients. *J NIH Res*. 1997;9:34-35
- Lee WM. Drug induced hepatotoxicity. *N Engl J Med*. 1995;333:1118-1127.
- Magil MK, et al. A multiple chemical sensitivity syndrome. *Am Acad Fam Phys*. [www.aafp.org/afp/980901ap/magil.html](http://www.aafp.org/afp/980901ap/magil.html).
- McKinnon RA, et al. Possible role of cytochromes P450 in lupus erythematosus and related disorders. *Lupus*. 1994;3:473-478.
- Meyer, et al. Genetic polymorphisms of drug metabolism. *Prog Liver Dis*. 1990; 9:307-323.
- Nassauto Get al. Effects of silibinin on biliary lipid composition. Experimental and clinical study. *J Hepatol* 1991 ;12:290-295.
- Oura H, Hiai S, Nabatini S, Nakagawa H, et al. Effect of ginseng on endoplasmic reticulum and ribosome. *Planta Medica* 1975;28:76-88.
- Oura H, Hiai S, Seno H. Synthesis and characterization of nuclear RNA induced by Radix ginseng extract in rat liver. *Chem Pharm Bull* 1971 ;19:1598-1605.
- Palasciano G, Protinacasa P, et al. The effect of silymarin on plasma levels of malonaldehyde in patients receiving long-term treatment of psychotropic drugs. *Curr Ther Res* 1994;55:537-545.
- Perlmutter D. Parkinson's diseases - new perspectives. *Townsend Lett*. Jan 1997: 48-50.
- Pointel JP, Boccalon H, Cloarec M, et al. Titrated extract of *Centella asiatica* (TECA) in the treatment of venous insufficiency of the lower limbs. *Angiology* 1987;38(11):46-50.
- Ridgen S, et al. Evaluation of the effect of modified entero-hepatic resuscitation program in chronic fatigue syndrome patients. *J Adv Med*. 1998;11(4):247-262.
- Ridgen S. Entero-hepatic resuscitation program for CFIDS. *The CFIDS Chronicle*. Spring 1995:46-49.
- Skvortsova RI, Pzniakovski VM, Agarkova IA. Role of vitamin factor in preventing phenol poisoning. *Vopr Pitan* 1981; 2: 32-35.
- Sonnenbicher J, Goldberg M, Hane L, et al. Stimulatory effect of silibinin on the DNA synthesis in partially hepatectomized rat livers. Nonresponsive in hepatoma and other malignant cell lines. *Biochem Pharm* 1986;35:538-541.
- Steventon GB, et al. Xenobiotic metabolism in Parkinson's disease. *Neurology*. 1989;39:883-887.
- Susnik F. Present state of knowledge of the medicinal plant *Taraxacum officinale*. *Weber. Med Razgledi* 1982;21:323-328.
- Wagner H. Antihepatotoxic flavonoids. In: Cody V, Middleton E, Harbourne JB, eds. *Plant flavonoids in biology and medicine: biochemical, pharmacological, and structure-activity relationships*. New York, NY: Alan R Liss. 1986:p545-558
- Yammato M, Uemura T, Nakama S, et al. Serum HDL-cholesterol-increasing and fatty liver-improving action of Panax ginseng in high cholesterol diet-fed rats with clinical effects on hyperlipidemia in man. *Am J Chin Med* 1983;11:96-101.
- Jinsheng Yu, Sharon Marsh, Junbo Hu, Wenke Feng, and Chaodong Wu, "The Pathogenesis of Nonalcoholic Fatty Liver Disease: Interplay between Diet, Gut Microbiota, and Genetic Background," *Gas-troenterology Research and Practice*, vol. 2016, Article ID 2862173, 13 pages, 2016. <https://doi.org/10.1155/2016/2862173>
- Chen, Yung-Ju et al. "Dietary broccoli protects against fatty liver development but not against progression of liver cancer in mice pretreated with diethylnitrosamine" *Journal of functional foods* vol. 24 (2016): 57-62.