

# ProBile+

Supports Healthy Gall Bladder Function



Biogenetix  
Nutrition Uncompromised

## PRODUCT BENEFITS

- Support and stimulate bile production and elimination
- Provides support for a healthy gallbladder
- Provides support for healthy bile flow

ProBile+ provides herbal compounds that support the production, flow, and elimination of bile. ProBile+ also works to support a healthy gallbladder.



## DIRECTIONS FOR USE

Take 1 capsule once per day, or as directed by your health care professional.

## STORAGE

Store in a cool, dry place away from direct sunlight. Product from unopened containers that have been stored under recommended conditions (55-85° F / 13-29° C) may be used for up to 36 months.

## WARNING

Consult your health care professional before use if pregnant, nursing, taking medications, or for any use by minors. Do not use if safety seal is broken or missing. Keep out of reach of children.

## SUPPLEMENT FACTS

|   | Amount<br>Per Serving | % Daily<br>Value |
|---|-----------------------|------------------|
| Serving size: 1 Capsule<br>Servings per container: 90 |                       |                  |
| Vitamin C (as ascorbic acid)                          | 36mg                  | 40%              |
| Dandelion Root Extract                                | 240 mg                | **               |
| Soy Bean Lecithin (40% Phosphatidylcholine)           | 72 mg                 | **               |
| Milk Thistle Seed Extract (80% Silymarin)             | 68 mg                 | **               |
| Ginger Root   | 55 mg                 | **               |
| L-Taurine   | 44 mg                 | **               |
| Turmeric Root Extract (95% Curcuminoids)              | 32 mg                 | **               |
| Beet Root   | 22 mg                 | **               |
| Cellulase   | 22 mg                 | **               |
| Protease  | 22 mg                 | **               |
| Lipase  | 22 mg                 | **               |
| Bitter Orange Fruit Extract (40% Hesperidin)          | 12 mg                 | **               |
| Sarsaparilla Root                                     | 4 mg                  | **               |
| Catalase (0.003 BU)                                   | 20 mcg                | **               |
| Super Oxide Dismutase (0.16 MFU)                      | 20 mcg                | **               |
| ** Daily Value not established.                       |                       |                  |

**Other Ingredients:** Hypromellose(Capsule),Magnesium Stearate, Silicon Dioxide, Microcrystalline Cellulose.

**Allergen:** Contains soy (as lecithin).

## REFERENCE LIST

1. Adzet T. Polyphenolic compounds with biological activity and pharmacological activity. *Herbs Spices Medicinal Plants* 1986;1:167-184
2. Bohm K. Choleretic action of some medicinal plants. *Arzneimittel Forsch* 1959;9:376-378.
3. Faber K. The dandelion *Taraxacum officinale*. *Pharmaize* 1958;13:423-436.
4. Feilbrich F, Koch H. Silymarin an inhibitor of lipoxygenase. *Experientia* 1979;35:148-150.
5. Fiebrich F, Koch H. Silymarin, an inhibitor of prostaglandin synthetase. *Experientia* 1979;35:150-152.
6. Giri J, Sakthi Devi TK, Meerarani S. Effect of ginger on serum cholesterol levels. *Ind J Nutr Diet* 1984;21 :433-436.
7. Gujral S, Bhumra H, Swaroop M. Effects of ginger (*Zingiber officinale* Roscoe) oleoresin on serum and hepatic cholesterol levels in cholesterol fed rats. *Nutr Rep Intl* 1978;17:183-189.
8. Hannin I, Ansell GB. *Lecithin. Technological, biological, and therapeutic aspects.* New York, NY:Plenum Press;1987.
9. Hikino H, Kiso Y, Wagner H. Antihepatotoxic actions of flavanolignans from *Silybum marianum* fruits. *Plant Medica* 1984;50:248-250.
10. Jenkins SA. Vitamin C and gallstone formation: a preliminary report. *Experientia* 1977;33:1616-1617.
11. Nassauto G et al. Effects of silibinin on biliary lipid composition. Experimental and clinical study. *J Hepatol* 1991;12:290-295.
12. Palasciano G, Protinacasa P, et al. The effect of silymarin on plasma levels of malonaldehyde in patients receiving long-term treatment of psychotropic drugs. *Curr Ther Res* 1994;55:537-545.
13. Sonnenbicher J, Goldberg M, Hane L, et al. Stimulatory effect of silibinin on the DNA synthesis in partially hepatectomized rat livers. Non-responsive in hepatoma and other malignant cell lines. *Biochem Pharm* 1986;35:538-541.
14. Srinivasan K, Sambaiah K. The effect of spices on cholesterol 7 alpha-hydroxylase activity and the serum and hepatic cholesterol levels in rat. *Int J Vitam Nutr Res* 1991 ;61 :364-369.
15. Susnik F. Present state of knowledge of the medicinal plant *Taraxacum officinale*. *Weber. Med Razgledi* 1982;21 :323-328.
16. Tushilin SA, Drieling DA, Narodetskaja RV, Lukash LK. The treatment of patients with gallstones by lecithin. *Am J Gastroenterol* 1976;65:231.
17. Wagner H. Antihepatotoxic flavonoids. In: Cody V, Middleton E, Harbourne JB, eds. *Plant flavonoids in biology and medicine: biochemical, pharmacological, and structure-activity relationships.* New York, NY: Alan R Liss. 1986;p545-558